METHOD OF CALCULATING SOIL TEMPERATURE AT A DEPTH OF 5 cm FROM ATMOSPHERIC TEMPERATURE AND HUMIDITY VALUES

L. N. Popovich

Translation of "K metodike rascheta temperatury pochvy na glubine 5 cm po znacheniyam temperatury i vlazhnosti vozdukha," Trudy Nauchno-Issledovatel'skogo Gidrometeoro-logicheskogo Instituta, Kiev, No. 91, 1970, pp. 89-98.

(NASA-TT-F-14516) METHOD OF CALCULATING
ATMOSPHERIC TEMPERATURE AT A DEPTH OF 5 CM FROM
L. V. Popovich (Translation Consultants,

CSCL 08M G3/13 36020

METHOD OF CALCULATING SOIL TEMPERATURE AT A DEPTH OF 5 cm FROM ATMOSPHERIC TEMPERATURE AND HUMIDITY VALUES

L. V. Popovich

ABSTRACT. The application of a method for soil temperature determination, previously proposed, is discussed, calculations are made, and graphs and charts are compiled for use in determining soil temperature at a depth of 5 cm under natural cover.

Study of the temperature regime in the arable layer of the soil, that in \(\sqrt{89*} \) which the main mass of plant roots is found, is of practical value for agricultural production. Intensity of growth, and plant development [1], definitely depend on soil temperature, particularly during the initial period of vegetation development. The thermal regime in the soil has a great influence on the rate at which biological and biochemical processes take place in the soil, and hence on soil fertility. Consequently, soil thermal conditions play an important part in the complex set of agrometeorological characteristics determining crop yields.

The possibility of determining the conditions of the soil temperature regime, averaged for a definite area, is of great practical importance, and something that cannot always be done by direct measurement. The proposed method of making the calculation provides the characteristics of the temperature regime, the average for meteorological areas covered by a natural plant cover (grassland) that is cut from time to time. The basis for this method of making the soil temperature determination was set forth in earlier articles by the author [3,44]. These papers pointed out that so-called inertia, that is, delay (shift in phases) in daily and seasonal air temperature variations at a height of 2 meters as compared to the variation in soil temperature at the depth considered, had to be taken into consideration in order to establish the quantitative connection between air temperature and soil temperature.

^{*} Numbers in the margin indicate pagination in the foreign text.

This paper applies the procedure set forth earlier for calculating the mean 10-day values of soil temperature at a depth of 5 cm at 0100, 0700, 1000, 1300, and 1900 hours.

First to be studied was the inertia in the daily variation in air temperature and humidity at a height of 2 meters, and in soil temperature at 5 cm. was done by a detailed plotting [2, 3] of curves for the daily variation in these elements from the mean 10-day values for six observation times (0100, 0700, 1000, 1300, 1600 and 1900 hours). These curves, even in their external appearance, show a delay in soil temperature maxima at 5 cm as compared to the maximum air temperatures at a height of 2 meters. The curves for daily variation in soil temperature at 5 cm were replotted at a different scale to bring their daily variation amplitudes into coincidence with the same amplitudes $\hat{\chi}$ in the air, so as to include the real delay in all the other corresponding temperature phases that can differ from the delay for the maxima at different times. The correction factors for daily inertia in heat transfer between air temperature at 2 meters and soil temperature at 5 cm were taken from these curves and plotted in terms of air temperature and humidity values. The result was an individual correction factor nomogram for each time of observation. relationships are presented at the end of this paper in the form of tables (Appendix I).

The delay in the onset of extremal temperatures at 5 cm, as compared togair temperature at 2 meters, is shorter than it is at the 10 and 20 cm levels, so the magnitudes of the correction factors for the depth were changed accordingly. Correction factors calculated for the 5 cm level were smaller in absolute values

for all times than were the same factors calculated for the 10 and 20 cm depths. Correction factor maxima (-2, -6°C) were a maximum in the morning hours (0700, 1000), at which time the maximum lag in the phase of soil temperature oscillations behind the corresponding phase in the air was observed.

The intensity in the heat transfer from the surface of the soil decreases, as does the difference between soil and air temperatures, during the second half of the day. The correction factors for 1300 and 1900, for example, decrease to $1.0 - 0.1^{\circ}$. The least correction factor $(0.1 - 0.5^{\circ})$ is observed at 0100 (Appendix I).




Figure 1. Magnitudes of daily correction factors for air water vapor pressure for 0100 hours.

Figure 1 shows the values for the magnitudes of the correction factors for humidity at 0100 hours. Air temperature is plotted on the x-axis, air water vapor pressure on the y-axis. As we see, the correction factor for this time ranges from -0.1 to 0.4 mb. The maximum error in determination of the water vapor pressure from the humidity tables is ±0.3 mb, if the dry and wet-bulb

temperature readings are accurate to within 0.1°. These correction factor magnitudes cannot, therefore, be taken into consideration. The exception is the 1000 observation, at which time isolines from -0.5 to 1.5 mb can be drawn. But even these correction factor magnitudes are but an insignificant part of the eigenvalue of the air water vapor pressure. It therefore is possible to ignore possible correction factors for inertia in the daily variation in air humidity. Determination of soil temperature at a depth of 5 cm then become much simpler than is the case for depths of 10 and 20 cm.




Figure 2. Curves from soil temperature calculations at 5 cm for individual months.

A series of curves were constructed (Figure 2) to calculate soil temperature. Air temperature at a height of 2 meters, corrected for daily inertia in heat transfer, was plotted on the x-axis, and uncorrected humidity values were plotted on the y-axis. Graphical relationships were constructed for each of the months \(\frac{9}{2} \) April through October, precluding the need to consider seasonal corrections for air temperature and humidity. Mean 10-day values of soil temperature, determined from Figure 2 and measured for the same times, were compared in order to evaluate the accuracy of the soil temperature calculation curves thus obtained. The comparison was made using data from the stations providing the observations used to construct the curves, but for a different period. Overall, 60 verifying calculations were made for each of the graphics in Figure 2. Absolute error distribution curves were approximately symmetrical, indicating an absence of

systematic errors. Root-mean-square deviations (σ) in calculated temperatures from measured temperatures also were calculated. σ was the same for each of the months: April 1.6°, May 1.3°, June 1.6°, July 1.6°, August 1.4°, September 2.1°.

Table 1 list the probability of different magnitudes of deviations in actual temperatures from those calculated from the curves in Figure 2.

TABLE 1.

TEMPERATURES VALUES FROM MEASURED VALUES Magnitude of deviation, Months 1,1-2,0 2,1-3,0 3,1-4,0 4,1-5,0

PROBABILITY OF DEVIATION (%) IN CALCULATED SOIL

April 44,4 23.3 38.3 63.3 41.6 11.6 53,3 60 3,3 6,6 25 38.3

Data provided by the Poltava meteorological station for the 0700 observations for May 1964, will be used as an example of the calculation. Air temperature and humidity values at a height of 2 meters, averaged for the first 10-day period, were 10.8° and 9.3° mb, respectively. Appendix I provides the air temperature correction factor $\Delta T = -2.7^{\circ}$. The corrected air temperature value $\sqrt{93}$ will be

$$T_{0}^{(i)} = T_{2m} + \Delta T = 10.8^{\circ} - 2.7^{\circ} = 8.1^{\circ}$$

Using the coordinates $T' = 8.1^{\circ}$ and $\hat{e} = 9.3$ mb in Figure 2 for May, we determine the soil temperature at a depth of 5 cm to be 10°. The actual soil temperature at the time was 9.3°.

The result of the processing was to obtain general graphical relationships, presented in Figure 2, for different soils and climatic zones, for use in finding the soil temperature at a depth of 5 cm under natural cover. The next stage of the work should be that of investigating the applicability of these curves to concrete forms of underlying surfaces. At the same time it should be emphasized that concretization here will mean seeking correction factors for the procedure developed, making this paper the basis of such future work.

Commas represent decimal points.

APPENDIX I. MAGNITUDES OF CORRECTION FACTORS (°C) FOR INERTIA IN THE DAILY VARIATION IN AIR TEMPERATURE

	9 20	4
	¥	4,0
	18	0000 4466
	17	000000 44460000
	16	00000000 446668111
	15	0000000 4466898111
	14	4466844111
qm	13	00,000,000,000,000,000,000,000,000,000
sure.	12	00100 4,0 0,0 0,0 1,0 1,0 1,0 1,0
Pressure	-	000000 mmdd-1-1
Vapor	10	
Water V	6	0,0000 E,E,0'4'-1'-1'-1'-1'-1'-1'-1'-1'-1'-1'-1'-1'-1'
Air Wa	.∞	600000 600000
A	2	00000 mmaa_
	တ	, , , , , , , , , , , , , , , , , , ,
	ທ	0,1
Air	temperature	0-2222222222222222222222222222222222222
•	te	

ŧ	A CONTRACTOR OF THE PROPERTY O		4.3)
l		٠.	• :	
				· :]
4.3				
			***	1
				·
	<u> </u>	_		
	ને વે વે વે વે વે			
		_		
	2222 & & & 444 d			3
	4444			
				··
	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4			
•				
		-		
	6 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4		N.	. :
				-
_	Odrodrod-rod-		7	
	૽ૺ૽૽૽ૺ૽ૺ૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽૽	. :	·	
				<u>.</u>
-	1 9 9 4 4 4 4 4 0 0 0 0 0 0 0 0 0 0 0 0 0			
	1971 1971 1971 1971 1971 1971 1971 1971	1		
:	: 17. [2] [2] [3] [4] [4] [4] [4] [4] [4] [4] [4] [4] [4	_		·
	-1900 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1.		
70	12222224444	1. 1		
ırs				<u></u>
on O		70		
h	7999997777	irs.		
0700 hour	-w04@00@04r0	hoúr		<u></u>
Ž	——∸ઇઇઇઇઇઇઇ ઇ ઇવ4440			•
0		2		٠.
	011244466664444 2460 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	1000		
		– (
		:	<u></u> ,	:
			٠.,	1,2
		- : ;		1
	0-1-244 6 6 6 4 4 4 6 7 0 0-70 6 7 0 4 8 - 7 7 0	5		F 00
	0112446664446 017067040-670	11.	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	-
				1 1
٠.	<u> </u>		- 6	2-
				~ C)
		• .	1:	
			∞ ~ ~	<u> </u>
			77	
				· · ·
			-2,0	• :
			T	
1		-		<u>. </u>
'			,	
1				
	0-0224222222222222222222222222222222222	2	∸ಇಣ	4. ro
1				
			-	

	20		77.70
	19		1111515 0.6.0.0.1.4.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0.0
	18		11144444646
	17		11119494466644 4000000000000000000000000000000
	16		1111199988888884444 οωναίτα ο ο ο ο ο ο ο ο ο ο ο ο ο ο ο ο ο ο ο
	15		
e, mb	4		1 1 1 1 1 2 2 2 2 2 2 2 2 2 2 2 4 4 4 4
Pressure	13	S	111114444444400 0 4 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2
1 1	12	1000 hours	11111444444444444444444444444444444444
Vapor	-	1000	
Water	10	,	11111111111111111111111111111111111111
ziń	6		1
 - -	8 0		044.0800000
	2		4 4 4 6 4 4 6
; -	. 9 :		80 8 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	[دن	,	
	re,		
Air	temperatur °C		\$5,800 0112E4757780828284888888888888888888888888888888

1	
	0 00000 1 00000 1 0 0 0 0 0 0
	00 00 00 00 00 00 00 00 00 00 00 00 00
	000 000 0-60000000000000000000000000000
	0,000 0,000,000,000,000,000,000,000,000
urs	0,000 0,000 1 0,1000 4 10 0 0 0 0 0
300 hours	66666 66666 666 666 666 666 666 666 66
13	
	00000 0000 0-000 0-000
	000000 000000 0-56640
	0,000
	0.0000
	0,00
	- 252 252 252 252 252 252 252 252 252 25
	AND

	20	
ssure, mb	19	10000 0000 0000 0000
	18	-000000000 -00000000000000000000000000
	17	1000000000 000000000 0000000000
	16	
	15	-000000000 -000000000
	14	-000000000 000000000
r Pre	13	1,1,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0,0
ater Vapor	12	900 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
		11 24-1-00000000000000000000000000000000000
Air W	10	441-1-1-0000
	6	211100000 211000000
	80	000000000000000000000000000000000000000
		1111100000
	9	
	ນ	<u>2</u>
	Temperature,	40.00.00012242572808282828

REFERENCES

- 1. Maksimov, N. A., <u>Kratkiy kurs fiziologii rasteniy</u> [Short Course in Plant Physiology], Sel'khozgiz, Moscow, 1958.
- Konstantinov, A. R., Goysa, N. I., Kudina, A. V., Levenko, A. A., "Calculation of the Delay in the Daily and Seasonal Variations in Air Temperature and Humidity at a Height of 2 Meters," <u>Trudy UkrNIGMI</u>, No. 41, 1964.
- 3. Popovich, L. V., "The Possibility of Calculating Soil Temperature at a Depth of 20 cm From Air Temperature and Humidity Data Measured at a Height of 2 Meters," Trudy UkrNIGMI, No. 78, 1969.
- 4. Popovich, L. V., "The Possibility of Calculating Soil Temperature at a Depth of 10 cm Under Grassland From Air Temperature and Humidity Values Measured at Meteorological Stations," <u>Trudy UkrNIGMI</u>, No. 87, 1969.

Translated for the National Aeronautics and Space Administration under contract NASw-2038 by Translation Consultants, Ltd., 944 So. Wakefield St., Arlington, Virginia 22204.